

Evaluating, Preparing and Amending **Lawn and Garden Soil**

Reviewed by Dave Franzen
Extension Soil Science Specialist

One of the biggest steps to establishing and growing turf, vegetables, ornamentals or flowers successfully is understanding the soil that provides their physical support and supplies them with water and nutrients.

Sometimes, the original soil will serve as an excellent growth medium, while other soils may need to be amended or even replaced with more adequate topsoil. This publication will help serve as a guide to evaluating soils and suggest amendments that will improve the success of lawn and garden plantings.

Soil evaluation should be conducted physically and chemically. Physical characteristics affect the ability of roots to explore the soil for water and nutrients, and the capacity of the soil to hold water as a reserve for plants between rains or irrigation. They also affect how easily water moves through the soil, preventing ponding.

While not a physical characteristic, the topography of the soil surface also will be an influence on water movement into the soil.

Physical Characteristic Terminology

Aeration – physically amending or altering soil to allow more oxygen to enter the soil and carbon dioxide to exit, thereby enhancing root health

Bulk density – the weight of dry soil per unit of volume. Well-aerated soil has a relatively low bulk density; compacted soil has a high bulk density.

Permeability – the ability to allow water to move freely through soil

Porosity – the percent of soil volume devoted to pore space. Compacted soils have lower pore space and higher bulk density.

Texture – the percent of sand, silt and clay that makes up the mineral portion of soil. Texture can be estimated by feeling for sand grains between your fingers and by the ability to produce a clay ribbon between your thumb and pointer finger (Figure 1). If necessary, texture can be measured accurately with a soil evaluation.

Fine textures – soil with more clay than sand or silt

Medium textures – soil with no dominance of sand, silt or clay

Coarse textures – soil with more sand than silt or clay

Water-holding capacity – the amount of water a soil can hold. Soil can hold two types: water that is plant-available and water that is held hydroscopically to soil particles and is unavailable for plant use. A silt loam has the most plant-available water but clay soil holds the most total water. Medium-textured soil holds more plant-available water than coarse-textured soil.



Figure 1. Determining clay content in soil via the ribbon test. This would be an example of a high-clay content soil.

Soils in nature develop in layers, called horizons (Figure 2). The sum of horizons from top to bottom is called the soil profile.

The uppermost layer of soil in most of North Dakota developed under



Figure 2. A slice of soil showing the various layers or horizons that have been formed through time.

prairie grasses and forbs, and tends to be dark-colored due to the resulting organic matter accumulation. This layer is the most productive in the soil.

The organic matter and biological activity in the zone helps soil particles clump together into aggregates. These aggregates tend to resist compaction and contribute greatly to permeability, favorable bulk density and porosity. The surface layer, rich in organic matter, also is a storehouse of slow-release nutrients that are helpful for plant growth.

The underlying layers are not as productive as the surface layer due to lower organic matter content. Sometimes the subsurface layers have been altered with an accumulation of soluble salts and carbonates. These accumulations generally are not desirable and may limit the cultivation of some types of plants.

With ornamentals, having a subsoil chemical analysis before selecting adapted species for planting is particularly important. Subsoils cannot be improved easily.

Soil Evaluation

Soil evaluation should begin with physical observations. Physical problems or limitations are most difficult to remedy, so knowing if the soil has any restrictions on use is important to determine.

Aeration

Evidence – Look for an abundance of fine, white roots under previous plant growth. Well-aggregated soils with bright yellow subsoil are indicators of well-aerated subsoils. Poorly aerated soils may have a musty smell from fungal growth or be dark, drab gray or olive.

Remedies

In general, poor aeration is caused by high clay content, compaction and/or natural depressional landscape positions. Take care to not overwater. Choose a better landscape position for garden plants or consider a raised bed.

Lawns: Mechanical aerators can be used to alleviate compaction. Natural shrinking and swelling of soil clay will fill in the cavities made by the coring machine, forming smaller aggregates.

Gardens: To amend clays, you should add sand only if using rates of 80 parts of sand to one part of soil. Additions of unmilled sphagnum peat moss, reed-sedge peat or hypnum peat are better amendments. The peat serves to separate soil particles, and the biological activity it promotes results in better aggregation of soil particles and separation of clays, making them more resistant to compaction.

Chemical Evaluation

A soil test is the best way to evaluate nutrient needs in plants, regardless of whether you plan to use an organic or chemical approach to nutrition. Home test kits provide general guidelines.

North Dakota recommendations are based on locally calibrated laboratory tests to get more reliable and repeatable results. Sending a soil sample to local labs across the state or to NDSU's soil testing lab on campus in Fargo will assure more accurate results with more reliable recommendations.

Collect samples from several locations around your lot (roughly one sample per 1,000 square feet), with each sample consisting of enough soil to fill a bag that can hold a sandwich. Different samples need to be collected for each location: backyard, front yard, etc.

Collect samples from 3 to 6 inches deep using a trowel, shovel or soil probe. Do not include the thatch layer with the soil. Mix the samples from each area to be tested separately using a clean bucket and send in 1 cup of soil following the lab's instructions for submitting samples.

Request a standard turfgrass test, which includes pH, soluble salts (EC), nitrogen (N), phosphorus (P), potassium (K) and organic matter content. While these three nutrients (N, P and K) are considered primary for good turfgrass growth, they are only three of the 13 mineral elements found in the soil that are necessary for plant growth.

Nitrogen is the key element in turfgrass growth, and the level of soil nitrogen can vary depending on the quality of the turf desired.

Minimally maintained lawns should receive a single application of fertilizer in the fall at a rate of 0.75 to 1 pound of actual N per 1,000 square feet;

high-quality lawns should receive two to three applications of fertilizer, once in the fall, again in the spring after at least one mowing of green grass and a third application around Memorial Day weekend.

The form nitrogen comes in is important; WSN (water-soluble nitrogen) quickly stimulates grass growth, causes increased mowing frequency and is short-lived.

When selecting fertilizer, look for at least one-third of the nitrogen source to come from WIN (water-insoluble nitrogen) sources for longer-lasting effect and not as high a mowing frequency.

The other two elements on the fertilizer bag, phosphorus and potassium, usually are not an issue in most home lawn situations. If a soil test indicates a serious deficiency of these two elements, recommendations will include specific formulations.

Existing lawns seldom will be deficient in P or K if the homeowner follows any kind of fertilization program and returns the grass clippings to the lawn most of the time. For new lawns, incorporating any needed P or K into the topsoil following lab recommendations is advisable before seeding or installing sod.

Garden Soils

The soil sample should be taken from the top 6 inches. In new gardens, a subsoil sample from the 6- to 18-inch depth for a salt analysis and soil pH is suggested.

When coring below 6 inches, call the One-Call 800 number in your area to avoid contacting or cutting utility cables. If these obstacles are in the intended garden area, consider another site for the garden.

If N levels are below the critical level, apply 4 to 5 pounds of 10-10-10

How to Read a Fertilizer Bag

All fertilizers registered for sale are required to be analyzed for their plant nutrient content and display the results on the bag: for example, 30-5-10. In this instance, the numbers refer to the amount of nitrogen, phosphorus pentoxide and potassium oxide, but simply are expressed as N, P and K. A 35-pound bag of fertilizer with this analysis would have 30 percent N (10.5 pounds of N), and 5 and 10 percent of the other two materials, respectively. Often the single element of nitrogen is sold as Ureaform, with an analysis of 46 percent nitrogen, and would show up on the bag as 46-0-0.

In North Dakota, the soil often is quite alkaline (high pH), resulting in the tie-up of a particular element, iron (Fe). When the usual fertilization practices don't green up the lawn to full potential, the problem may be an iron deficiency, characterized by a yellowing (chlorosis) of the turf.

In that case, look for a fourth number on the bag following potassium to indicate iron content, generally 3 to 5 percent, and usually combined with sulfur (S) as iron sulfate.

Although Earth has more than 100 elements, only 14 mineral nutrients are required universally for plants to grow and produce fruit or seed. They are listed below with their chemical symbols:

nitrogen (N), phosphorus (P), potassium (K), sulfur (S), calcium (Ca), magnesium (Mg), zinc (Zn), copper (Cu), manganese (Mn), boron (B), molybdenum (Mo), iron (Fe), chloride (Cl), nickel (Ni)

(or something close to this analysis) per 1,000 square feet. Seasonal applications of nutrients can be made using a highly soluble complete fertilizer (for example, Miracle-Gro or comparable fertilizer) when watering after the plants are established.

Organic Gardens

A soil sample should be used to guide amendments. Generally, organic gardening practices with the heavy use of plant and animal waste from composting are not in need of any drastic additions. The soil nutrient levels are usually in good balance for optimal plant growth.

Sources such as Organic Gardening magazine, NDSU's website at www.ag.ndsu.edu/pubs/plantsci/hortcrop/h1106.pdf or Ohio State's site at <http://ohioline.osu.edu/hyg-fact/1000/1257.html> will provide useful information in guiding the gardener who wishes to follow organic practices.

Crop rotation and cover crops are common practices with organic gardeners. As the growing season comes to an end, sow an oat or rye crop when the seasonal vegetables have been removed to help build the soil and mitigate the common home gardener habit of planting the same vegetables in the same spot each year.

Compaction

Compaction reduces soil porosity, which means air and water have more trouble penetrating and moving through the soil. Air movement and root penetration are restricted in compacted soils. Roots move readily in larger pores, but may not have enough strength to part soil particles between smaller pores.

Evidence – Water runs off rather than moving into the soil. Water may pond rather than infiltrate into the soil.

Roots tend to be restricted to the top few inches. In lawns, rooting is mostly in the top couple of inches. Roots may grow sideways rather than down.

Remedies – Lawns: Mechanical aerators result in better aggregation and aeration. Limit traffic when soil is wet.

Gardens: Till deeply but with care not to bring up undesirable soil containing salts or carbonate minerals. Add generous amounts of peat moss or compost.

High Soil Sodium

Spots in the yard or garden where water ponds may indicate areas of sodium accumulation. Soils dry hard and, when watered, water tends to pond very quickly. Sodium prevents soil particles from aggregating, instead forming large monoliths with soil particles so tightly bound that roots can explore only the surface. Plants growing in these areas require frequent watering and grow poorly, if at all.

Evidence – You'll see areas of poor plant growth, ponding and soils that are hard to till. Often the surface of the soil has a white crust, and a soil test of the top 6 inches shows high levels of sodium present.

Remedies – Better drainage is required for any amendment to be effective. A soil test can provide an SAR, or sodium absorption ratio, showing the relative balance of sodium to the desirable cations of calcium and magnesium.

The higher the SAR reading, the greater amount of soluble calcium amendments you'll need to replace sodium on soil clays and flush them away with rainfall or high-quality (low-sodium) irrigation water. Amounts required may range from 20 to 200 pounds per 1,000 square

feet of gypsum. The amendments will need to be tilled in to the depth of the sodium problem.

Several rooting depths of water may be required to leach out the sodium. If the soil has limited drainage and the irrigation water is high in sodium salt, the probability of positive results using this technique is slim.

Growing grasses that are tolerant to drought, considering moving the garden or building raised beds that have better drainage and contain soil that is not degraded by excess sodium would be better.

Layering

Plants develop better in deep, homogenous soil layers that are similar in texture. A few inches of good topsoil over some not-so-good soil will result in plant roots growing mostly in just the good soil. This restricts the amount of soil used for water and nutrient uptake, which will require more frequent watering and fertilization.

Evaluation – Soil core samples indicate soil textural changes in the common rooting zone. The rooting zone for lawns is about 6 inches. Most garden plants need about 18 inches.

Remedies – Lawns: Before seeding, make sure that the topsoil is at least 6 inches deep; deeper is better.

Gardens: Amendments need to be tilled in deeply. Avoid tilling in organic amendments or sand only to the surface.

Raised beds need to be as uniform as possible to the entire depth. If raised beds are shallower than 18 inches, incorporating the original soil – if not contaminated with salts or undesirable materials – into the added soil material for raising will increase effective rooting.

Salts

Salty soils are a natural, but undesirable, result of relatively young soils. They are high in nutrient content, have poor drainage and are often in a semiarid climate (Figure 3). Some plants have a higher salt tolerance than others. Most garden plants and many desirable lawn grasses have a low tolerance for salt.

When the water table is high (the water table is a fluctuating zone under the soil where the soil is saturated with groundwater for a long period of time), salts are brought to the surface by capillary action. The water is pulled toward the soil surface and evaporates, leaving the salts behind.

During times of continuous dry weather, any rainfall received tends to drive the salts deeper. In wetter periods, salty areas expand as water tables move closer to the soil surface.

Evaluation – A sign of salty soils is white, chalky material at the soil surface. Plants growing in or near these areas may be stunted or dull, have scorched-looking leaves or leaf margins and require more frequent watering. Soil tests indicate EC (electrical conductivity) is greater than 0.5 millimoh per centimeter (or decisiemen/meter).

Remedies – No amendment that “neutralizes” salty soil is available.



Figure 3.
Salt-affected soil.
In a soil such as this,
growth of most horti-
cultural plants would
be very poor,
if at all.

Salts are present due to poor drainage and high water tables. Adding any amendment usually increases salt levels.

Test irrigation water for salts. If salt levels are high for your soil (NDSU Extension publications EB68 and AE792), a leaching irrigation after the season (for example, turfgrass, 6 inches; vegetables and flowers, 12 to 18 inches; trees and shrubs, 18 to 24 inches) might be needed to sustain productivity.

Lawns: If tiling is an option, you might consider it, especially in golf courses and natural athletic fields. Having an adequate outlet is often a problem and is a reason tiling is not used more often.

Before tiling, make sure that sodium will not be a problem if the salts are drained out of the soil. Usually salts only affect a certain area of a lawn. These areas might be better off seeded to a salt-tolerant grass (NDSU Extension publication SF1087, “Managing Saline Soils in North Dakota”).

Gardens: Consider moving the garden to a less salty spot if possible. If this is undesirable, consider a substantially raised bed at least 18 inches or taller. Fill the bed with new, low-salt soil and avoid incorporating the original soil into the bed.

Soil Cracking

Soil cracking is common in dry soils with high clay content. These cracks accelerate drying of subsoil and limit soil water-holding capacity during hot summer months.

Remedies – Lawns: Use more frequent, but less intense, waterings; maintain good grass coverage.

Gardens: Add organic amendments, especially unmilled peat moss, to improve aggregation. Do not overwater, but do not allow the soil to become overly dry.

Soil Crusting

Crusts may form following a hard rain or intensive irrigation on bare soil. Crusts are most harmful when establishing a lawn from seed or when a small-seeded crop, such as lettuce or carrots, has just been seeded into the garden.

Evidence – You can observe a hard surface layer after a heavy rainfall or when the soil dries.

Remedies – Lawn: Use a mulch to slow soil drying. Irrigate with smaller, less intense droplets. Gypsum may help reduce crusting in some soils.

Garden: If you noticed soil cracking in previous gardening years, amend soil with unmilled peat moss. A gypsum application may be helpful in some soils. Apply gypsum at rates of about 45 to 90 pounds per 1,000 square feet. Applications are recommended in the fall after crop removal to allow the chemical reactions to begin taking place. Gypsum generally is not as beneficial as organic amendments.

Soil Temperature

Around a home, soil temperatures can vary as much as 10 F at any given time. The north side of a house shades the soil for as much as two spring months more than the south side.

A thermometer can guide a gardener toward plants more adapted to cooler or warmer soil environments.

If the daytime temperature is too hot for some plants, mulching the soil after emergence or planting will help reduce soil temperature fluctuations and produce a cooler soil on average.

If soils are too cool, warming them is difficult. However, you can select plants that are desirable, yet require cooler temperatures and shade to thrive.

Texture

Take a small amount of soil in your palm and wet it slightly. Roll it into a small ball in your palm. Next, try to make a flat ribbon between your thumb and your first two fingers (Figure 1). Push the soil into your fingers with your thumb.

If you can make something that looks like a ribbon about one-quarter inch thick, the soil is at least medium-textured and may be fine-textured. If the ribbon breaks off readily before it reaches $\frac{3}{4}$ inch in length, the soil is medium-textured.

If you can make a ribbon longer than that, the soil is fine-textured. If the soil doesn't make a good ribbon and you can feel the sand grains easily, it is coarse-textured.

Thatch

Thatch is a surface layer of undecomposed organic material in established lawns that can be good for the turf-grass if it is $\frac{1}{2}$ inch or less in thickness.

If it is more than $\frac{1}{2}$ inch thick, it becomes restrictive to water, air and nutrient movement into the root zone. Excess thatch can develop hydrophobic properties that repel water. It also can harbor insects and diseases, and because of the restrictions of air and water movement, it limits rooting.

Evaluation – You'll notice excessive cushioning when walking on the turf. Sod appears loose due to lack of rooting. The response to fertilizer, water and pesticide applications is poor.

Using a knife or soil probe, cut into the sod and measure thickness; any thatch that is deeper than $\frac{1}{2}$ inch will cause problems.

Remedies – Thatch can be controlled through regular core aeration; power-raking; adjusting cultural practices, including fertilization, mowing and pesticide use; and selecting nonaggressive turf cultivars. It is not caused by clippings being returned to the surface when mowing.

Undesirable Minerals

In certain parts of North Dakota, primarily along the northern tier of counties, the shale bedrock that underlies most of the state is fairly close to the surface. In these areas, glaciation mixed the shale more abundantly in the soil. These shales often contain relatively high levels of cadmium, selenium and arsenic.

If in the course of tillage, small pieces of gray, flat rock (shale) are evident in the soil, the chances are greater than normal that higher levels of these undesirable minerals will find their way into leafy, green garden plants.

These elements tend to accumulate in green, leafy vegetables, particularly broccoli, cauliflower, brussels sprouts, kale and Swiss chard. They are

much less concentrated in grain and fruit crops such as sweet corn and tomatoes.

If you are in a region prone to high levels of undesirable minerals, growing leafy crops in raised beds with imported soil would reduce exposure to these minerals.

Also, some fertilizer amendments contain relatively high levels of undesirable mineral. The guaranteed analysis is required to list the nitrogen, phosphorus and potassium levels, as well as any other plant nutrients sold in the material.

Other "inert" ingredients are not listed. However, several states, including Washington, require fertilizer sellers of all products in the state to have a total analysis and list these products on their fertilizer analysis website.

In North Dakota, we have no such law, and some amendments sold are very high in some of these minerals, particularly those containing iron. A gardener can obtain a better idea of a fertilizer's contents by going to the state of Washington's website and others, or asking the company directly for a detailed analysis.

Water-holding Capacity

The water-holding capacity is the amount of plant-available water possible if the soil is wetted to a point called field capacity – approximately one-third of atmosphere suction. This is almost totally dependent on soil texture, but the amount of aggregation and organic matter extends a soil's capacity, while sodium decreases it.

- Coarse-textured soils –
1 inch per foot in depth
- Medium-textured soils –
about 1½ to 2 inches per foot in depth
- Fine-textured soils –
2½ inches per foot in depth

Checklist for Soil Evaluation – Garden Soil

Site Location _____

Date _____ Plant Species _____

Texture		Water-holding capacity
Coarse	<input type="checkbox"/>	1 inch/foot
Medium	<input type="checkbox"/>	2 inches/foot
Fine	<input type="checkbox"/>	2½ inches/foot
Aeration		Remedy
Excellent	<input type="checkbox"/>	None
Good	<input type="checkbox"/>	Peat moss additions
Poor	<input type="checkbox"/>	Tillage, peat moss additions, double digging
Compaction		Remedy
No	<input type="checkbox"/>	None
Yes	<input type="checkbox"/>	Peat moss additions, deep tillage
Soil cracking		Remedy
No	<input type="checkbox"/>	None
Yes	<input type="checkbox"/>	Avoid deep soil drying
Crusting		Remedy
No	<input type="checkbox"/>	None
Yes	<input type="checkbox"/>	Peat moss, low-intensity irrigation
Layering		Remedy
No	<input type="checkbox"/>	None
Yes	<input type="checkbox"/>	Thorough mixing of soil
Salts		Remedy
No	<input type="checkbox"/>	None
Yes	<input type="checkbox"/>	Improve drainage, raised bed, move garden
Sodium		Remedy
No	<input type="checkbox"/>	None
Yes	<input type="checkbox"/>	Improve drainage, apply gypsum if drainage and deep tillage is possible. Move garden; investigate water source
Undesirable minerals		Remedy
No	<input type="checkbox"/>	None
Yes	<input type="checkbox"/>	Raised bed, bring in alternative soil, avoid leafy and Cole family plants
Soil test evaluation		Remedy
Needs N	<input type="checkbox"/>	4 to 5 lbs N as 10-10-10 or equivalent available N Fertilizer
Needs P	<input type="checkbox"/>	4 to 5 lbs P ₂ O ₅ as 10-10-10 or equivalent P source
Needs K	<input type="checkbox"/>	4 to 5 lbs K ₂ O as 10-10-10 or equivalent K source
Acid pH	<input type="checkbox"/>	Apply lime to achieve desired pH – about 6 lbs/400 SF
pH > 7	<input type="checkbox"/>	Plant adapted species, incorporate sphagnum peat
Organic Matter <3%	<input type="checkbox"/>	Add compost and/or peat moss and mix thoroughly
Soil EC > 0.5	<input type="checkbox"/>	Improve drainage; raised beds; plant salt-tolerant plants or move garden to lower-salt area

Checklist for Soil Evaluation – Lawns

Site Location _____

Date _____ Turfgrass Species _____

Texture		Water-holding Capacity
Coarse	<input type="checkbox"/>	1 inch/foot
Medium	<input type="checkbox"/>	2 inches/foot
Fine	<input type="checkbox"/>	2 ½ inches/foot
Aeration		Remedy
Excellent	<input type="checkbox"/>	None
Good	<input type="checkbox"/>	None
Poor	<input type="checkbox"/>	Mechanical core aeration
Compaction		Remedy
No	<input type="checkbox"/>	None
Yes	<input type="checkbox"/>	Mechanical core aeration
Soil cracking		Remedy
No	<input type="checkbox"/>	None
Yes	<input type="checkbox"/>	More frequent, but lower irrigation rates
Crusting		Remedy
No	<input type="checkbox"/>	None
Yes	<input type="checkbox"/>	Mulch new seedlings (straw, virgin wood fibers)
Layering		Remedy
No	<input type="checkbox"/>	None
Yes	<input type="checkbox"/>	Start lawn with at least 6 inches of good topsoil
Salts		Remedy
No	<input type="checkbox"/>	None
Yes	<input type="checkbox"/>	Improve drainage, seed-tolerant grass species
Sodium		Remedy
No	<input type="checkbox"/>	None
Yes	<input type="checkbox"/>	Improve drainage; apply gypsum if drainage and deep tillage is possible; seed drought-tolerant grasses; investigate or seek new water source
Thatch		Remedy
No	<input type="checkbox"/>	None
Yes	<input type="checkbox"/>	Power raking, core aeration; adjust cultural practices
Soil test evaluation		Remedy
Needs N	<input type="checkbox"/>	For normal lawns, clippings returned to turfgrass – 1 lb N/1,000 sq. ft. – late August
		Exceptional lawns, clippings removed – 2 to 3 lb total per season, distributed mid-May, late June and late August
Needs P	<input type="checkbox"/>	4 to 5 lb P ₂ O ₅ per 1,000 sq. ft. Do not exceed N recommendations
Needs K	<input type="checkbox"/>	4 to 5 lb K ₂ O per 1,000 sq. ft. Do not exceed N recommendations.
pH > 7	<input type="checkbox"/>	Plant adapted species
Soil EC > 0.5	<input type="checkbox"/>	Improve drainage; plant adapted cultivars

Some sources with useful information for gardeners who want to follow organic practices are:

- Organic Gardening magazine
- NDSU's website at
www.ag.ndsu.edu/pubs/plantsci/hortcrop/h1106.pdf
- Ohio State University's website at
<http://ohioline.osu.edu/hyg-fact/1000/1257.html>

This publication was authored by Ron Smith, retired;
and Dave Franzen, Extension Soil Science Specialist

For more information on this and other topics, see www.ag.ndsu.edu

NDSU encourages you to use and share this content, but please do so under the conditions of our Creative Commons license. You may copy, distribute, transmit and adapt this work as long as you give full attribution, don't use the work for commercial purposes and share your resulting work similarly. For more information, visit www.ag.ndsu.edu/agcomm/creative-commons.

North Dakota State University does not discriminate on the basis of age, color, disability, gender expression/identity, genetic information, marital status, national origin, public assistance status, sex, sexual orientation, status as a U.S. veteran, race or religion. Direct inquiries to the Vice President for Equity, Diversity and Global Outreach, 205 Old Main, (701) 231-7708.

County Commissions, NDSU and U.S. Department of Agriculture Cooperating.
This publication will be made available in alternative formats for people with disabilities upon request, (701) 231-7881.

1.5M-3-07; 1M-4-09; 500-7-11; 1M-3-13; 500-3-14